



North Carolina Department of Health and Human Services
Division of Public Health

Pat McCrory
Governor

Aldona Z. Wos, M.D.
Ambassador (Ret.)
Secretary DHHS

Penelope Slade-Sawyer
Division Director

Date: September 9, 2014, (2 pages)

To: All North Carolina Health Care Providers
From: Megan Davies, MD, State Epidemiologist
RE: **Respiratory Infections due to Enterovirus D68 (EV-D68)**

Enterovirus D68 (EV-D68) has recently been identified in association with respiratory illness outbreaks in the Midwest. As of September 9, 2014, no cases of EV-D68 infection have been confirmed in North Carolina. This memo is intended to provide general information regarding EV-D68 and recommendations for North Carolina health care providers.

Enteroviruses – Background

Enteroviruses are very common viruses. There are more than 100 types of enteroviruses. It is estimated that 10–15 million enterovirus infections occur in the United States each year. Most people infected with enteroviruses have no symptoms or only mild symptoms, but some infections can be serious. Most enterovirus infections in the United States occur seasonally during the summer and fall, and outbreaks of tend to occur in several-year cycles.

Clinical and Epidemiologic Features

EV-D68 is an enterovirus that was first isolated in California in 1962 and has been reported infrequently since that time. EV-D68 has been associated almost exclusively with respiratory disease, which can range from mild to severe. The full clinical spectrum of EV-D68 illness is not well-defined.

EV-D68 has been identified with increasing frequency during recent years, sometimes in association with large respiratory illness clusters in the United States and elsewhere. Whether this increase in recognized cases is attributable to improved diagnostics or emergence of the pathogen is unknown. EV-D68 infections appear to be more common in the fall. No data are available regarding the overall burden of morbidity or mortality from EV-D68 in North Carolina or in the United States.

Diagnosis

Enteroviruses can be detected by several commercial, multi-pathogen detection systems. However, the sensitivity of these systems for EV-D68 detection is unknown. Some of these systems use broadly reactive primers that amplify RNA from either human rhinoviruses or enteroviruses, and results are reported as "entero-rhinovirus" or "human rhinovirus/enterovirus". Most hospital and commercial laboratories are not able to perform enterovirus typing.

The North Carolina State Laboratory of Public Health (NCSLPH) offers viral culture to detect rhinoviruses and enteroviruses and serotyping for the detection of several specific enteroviruses, including enterovirus 70 and 71, echoviruses and coxsackieviruses.

www.ncdhhs.gov • <http://epi.publichealth.nc.gov/cd/>
Tel 919-733-7301 • Fax 919-733-1020

Location: 225 N. McDowell Street • Raleigh, NC 27603
Mailing Address: 1902 Mail Service Center • Raleigh, NC 27699-1902
An Equal Opportunity / Affirmative Action Employer



The gold standard test for EV-D68 identification is partial sequencing of the structural protein genes, VP4-VP2 or VP1. Facilities or providers interested in the specific detection of EV-D68 can submit specimens to the NCSLPH for submission to the CDC. Testing prioritization will be based on severity of illness. **Providers should contact their local health department or the Communicable Disease Branch epidemiologist on-call (919-733-3419) prior to submitting specimens.**

Acceptable specimen types include nasopharyngeal (NP) or oropharyngeal (OP) swabs in viral transport media (> 1ml) or an NP/OP wash or aspirate (> 1ml). Specimens should be collected within 1 week of illness onset, preferably within 48 hours post-onset. Specimens should be shipped on cold packs or dry ice. All specimen submissions must be accompanied by a completed CDC 50.34 DASH form (<http://slph.state.nc.us/Forms/CDC-Dash-NCSLPH-013114.pdf>), a NCSLPH virology form (<http://slph.state.nc.us/Forms/DHHS-3431-Virology-20130809.pdf>) and a descriptive case history for prioritization. Please contact Myra Brinson (919-807-8835) or Peggy Brantley (919-807-8820) at the NCSLPH if you have questions regarding enterovirus testing and specimen collection and shipment.

Treatment

There is no specific treatment for EV-D68 infections. Many infections will be mild and self-limited, requiring only symptomatic treatment. Patients with asthma exacerbations or other more severe manifestations may require hospitalization for supportive therapy. Vaccines for preventing EV-D68 infections are not currently available.

Public Health Recommendations

To help reduce the risk of infection with EV-D68 and other respiratory viruses, health care providers should recommend the following:

1. Wash hands often with soap and water for 20 seconds;
2. Avoid touching eyes, nose, and mouth with unwashed hands;
3. Avoid kissing, hugging, and sharing cups or eating utensils with people who are sick;
4. Disinfect frequently touched surfaces, such as toys and doorknobs, especially if someone is sick; and
5. Stay home when feeling sick, and obtain consultation from your health care provider.

Infection Prevention

Contact and Droplet isolation precautions are recommended for patients in whom infection with EV-D68 or another respiratory virus is suspected. Use of antimicrobial soap and water is preferred for hand hygiene, as alcohol-based hand rubs have limited activity against EV-D68 and other enteroviruses.

Reporting

Providers should be aware of EV-D68 as one of many causes of respiratory illness. Providers who identify clusters of severe or unusual respiratory illness are asked to contact their local health department or the Communicable Disease Branch epidemiologist on-call 24/7 number (919-733-3419).

Additional information is available at <http://www.cdc.gov/non-polio-enterovirus/about/EV-D68.html>. This is an evolving situation and recommendations may change as new information becomes available.

References

1. CDC (2014). Severe Respiratory Illness Associated with Enterovirus D68 — Missouri and Illinois, 2014. MMWR Morb Mortal Wkly Rep 63 – early release. Available at <http://www.cdc.gov/mmwr/pdf/wk/mm63e0908.pdf>. Accessed September 8, 2014.
2. CDC (2011). Clusters of acute respiratory illness associated with human enterovirus 68 – Asia, Europe, and United States, 2008–2010. MMWR Morb Mortal Wkly Rep 60, 1301–1304. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/mm6038a1.htm>. Accessed September 8, 2014.